

517.524

[dmytro\\_bodnar@hotmail.com](mailto:dmytro_bodnar@hotmail.com), [maria.bubnyak@gmail.com](mailto:maria.bubnyak@gmail.com)

1-

1-

1-

$$1 + D \frac{a_n}{b_n} = 1 + \frac{a_1}{b_1 + \frac{a_2}{b_2 + \dots}}, \quad (1)$$

$a_n, b_n (n \geq 1)$  ó ,  $k$ - ,  $a_{kn+p} = a_p$ ,  
 $b_{kn+p} = b_p (n = 0, 1, \dots; p = \overline{1, k})$ .

XIX . E. Galios, T. Thiele, A. Pringsheim. O. Perron [12],  
 H. Wall [13], W. Jones, W. Thron [7].  
 1966 . ( )

$$1 + D \sum_{k=1}^{\infty} \frac{a_{i(k)}}{b_{i(k)}} = 1 + \sum_{i_1=1}^N \frac{a_{i(1)}}{b_{i(1)} + \sum_{i_2=1}^N \frac{a_{i(2)}}{b_{i(2)} + \dots}}, \quad (2)$$

$a_{i(k)}, b_{i(k)}$  ó ,  $i(k) = i_1, i_2, \dots, i_k$  ó ,  $1 \leq i_k \leq N, k \geq 1, N$  ó  
 [10].

[6], [9], [8] [5],

$$1 + D \sum_{k=1}^{\infty} \frac{a_{i(k)}}{1} = 1 + \sum_{i_1=1}^{i_0} \frac{a_{i(1)}}{1 + \sum_{i_2=1}^{i_1} \frac{a_{i(2)}}{1 + \dots}}, \quad (3)$$

$a_{i(k)}$  ó ,  $i(k) = i_1, i_2, \dots, i_k, 1 \leq i_k \leq i_{k-1}, k \geq 1, i_0 = N$ .  
 (3)  $\emptyset$  [2], [3,4].

$$\left(1 + D \sum_{k=1}^{\infty} \frac{c_{i_k}}{1}\right)^{-1}, \quad (4)$$

$c_{i_k} \neq 0$  ó ,  $1 \leq i_k \leq i_{k-1}, k \geq 1, i_0 = N$ .

(3)

**I.**  $a_{i(k)}$  (3)

$$P_{i(k),\varepsilon}(\gamma) = P_{i_k,\varepsilon}(\gamma) = \left\{ \omega \in C : |\omega| - \operatorname{Re}(\omega e^{-i\gamma}) \leq 2p_{i_k} (1 - \varepsilon) \cos^2 \gamma/2 \right\},$$

$-\pi < \gamma < \pi, \varepsilon \acute{o}$  , (  $0 < \varepsilon < 1$  ),  $1 \leq i_k \leq i_{k-1}, i_0 = N$

$$p_{i_k} = \frac{1}{2N} \left( 1 - \frac{i_k}{2N} \right). \quad (5)$$

- 1) ;
- 2) (3) , :
- a)  $k$  ,  $a_{i(k)} = 0$  ( $i_p = \overline{1, i_{p-1}}, p = \overline{1, k}$ ),
- b)  $\sum_{k=1}^{\infty} \min \left( |a_{i(k)}|^{-1}; i_p = \overline{1, i_{p-1}}, p = \overline{1, k} \right)$  ,  $i(k)$ ,
- $a_{i(k)} = 0$  ;
- 3)

$$K(\gamma) = \left\{ z \in C : \left| z - \frac{e^{-i\gamma/2}}{\cos \gamma/2} \right| \leq \frac{1}{\cos \gamma/2} \right\}. \quad (6)$$

3.23 [6, .114]

(2).

$$V_{i(k)} = V = \left\{ \omega \in C : \operatorname{Re}(\omega e^{-i\gamma/2}) \geq -\frac{1}{2N} \cos \gamma/2 \right\}$$

$$E_{i(k)} = E_{i_k} = \left\{ \omega \in C : |\omega| - \operatorname{Re}(\omega e^{-i\gamma}) \leq 2p_{i_k} \cos^2 \gamma/2 \right\},$$

$1 \leq i_k \leq i_{k-1}, i_0 = N, k \geq 1, -\pi < \gamma < \pi, p_{i_k}$

(5).

$$(4) \quad 1 \quad (4)$$

**2.**

$c_{i_k} \in P_{i_k}(\gamma)$ ,

$$P_{i_k}(\gamma) = \left\{ \omega \in C : |\omega| - \operatorname{Re}(\omega e^{-i\gamma}) \leq 2p_{i_k} \cos^2 \gamma/2 \right\},$$

$-\pi < \gamma < \pi, i_k = \overline{1, N}, p_{i_k}$  (5).

- 1) (4) ;
- 2) (4)  $K(\gamma)$ ,
- (6).

$$\left(1 + D \sum_{k=1}^{\infty} \frac{z_{i_k}}{1}\right)^{-1}. \quad (7)$$

$$z_{i_k} \in P_{i_k}(\gamma) \quad (i_k = \overline{1, N}). \quad (7)$$

$$\sum_{k=1}^{\infty} \min(|z_i|^{-1}; i = \overline{1, N}) \delta$$

(7) ó

$$P_{i_k}(\gamma) \quad (i_k = \overline{1, N}),$$

$$\{F_n(z)\}_{n=1}^{\infty} \quad (7)$$

[6, .66],

$$\Delta = P_{i_k, \varepsilon}(\gamma).$$

$$z_{i_k} = c_{i_k}, \quad (4)$$

(4)

$$K(\gamma)$$

1.

I.  $a \neq 0$  ó

$$a \in E(p),$$

$$E(p) = \{z \in \mathbb{C} : |\arg(z+p)| < \pi\}, \quad p \text{ ó}$$

$$P(\gamma) = \{\omega \in \mathbb{C} : |\omega| - \operatorname{Re}(\omega e^{-i\gamma}) \leq 2p \cos^2 \gamma/2\}, \quad -\pi < \gamma < \pi. \quad (8)$$

$$1) \quad \arg a = \pi, \quad a \in P(\gamma) \quad - \quad \gamma;$$

$$2) \quad \arg a \neq \pi, \quad a \in P(\gamma), \quad \gamma_1 \leq \gamma \leq \gamma_2,$$

$$\gamma_1 = \begin{cases} \arccos\left(\frac{A+B}{C}\right), & \cos \alpha \leq 1 - \frac{2p}{|a|} \\ -\arccos\left(\frac{A+B}{C}\right), & \cos \alpha > 1 - \frac{2p}{|a|} \end{cases}, \quad (9)$$

$$\gamma_2 = \arccos\left(\frac{A-B}{C}\right) \quad (10)$$

$$\alpha = \arg a, \quad A = (|a| - p)(p + |a| \cos \alpha), \quad B = |a| \sin \alpha \sqrt{2|a|p(1 + \cos \alpha)}, \quad C = p^2 + |a|^2 + 2|a|p \cos \alpha.$$

$$a = |a|e^{i\alpha}, \quad a \in P(\gamma),$$

$$|a| - \operatorname{Re}(ae^{i(\alpha-\gamma)}) \leq p(1 + \cos \gamma).$$

$$(p + |a| \cos \alpha) \cos \gamma + |a| \sin \alpha \sin \gamma \leq |a| - p.$$

ø

$\gamma,$

:

$$\left| \gamma - \arccos \frac{p + |a| \cos \alpha}{\sqrt{C}} \right| \leq \arccos \frac{|a| - p}{\sqrt{C}},$$

$$\gamma \in [\gamma_1, \gamma_2], \quad \gamma_1, \gamma_2 \quad (9)-$$

(10).

$$-\pi < \gamma_1 \leq \gamma_2 < \pi.$$

$$G_1 = \{\omega \in \mathbb{C} : |\arg(\omega + p)| < \pi\} \quad (11)$$

$$c_1 \in G_1(\gamma) \text{ ó}$$

$$P_k(\gamma) = \{\omega \in \mathbb{C} : |\omega| - \operatorname{Re}(\omega e^{-i\gamma}) \leq 2p_k \cos^2 \gamma/2\},$$

$$p_k = \frac{1}{2N} \left(1 - \frac{k}{2N}\right), \quad -\pi < \gamma < \pi, \quad k = \overline{1, N}.$$

$G_s$

$$c_s \quad (s = \overline{2, N})$$

$$(4) \quad , \quad c_r \quad (r = \overline{1, s-1})$$

$$1 \quad : \quad c_1 \in P_1(\gamma) \quad , \quad \gamma \in [\gamma_1^1, \gamma_2^1], \quad \gamma_1^1 = \gamma_1 \quad \gamma_2^1 = \gamma_2,$$

$$(9)- \quad (10) \quad a = c_1 \quad p = p_1.$$

$$G_2 = G_2(c_1) = \bigcup_{\gamma \in [\gamma_1^1, \gamma_2^1]} P_2(\gamma).$$

$$c_2 \in G_2.$$

$$c_3 \cdot c_2 \in P_2(\gamma), \quad \gamma \in [\gamma_1^2, \gamma_2^2],$$

$$\gamma_1^2 = \gamma_1 \quad \gamma_2^2 = \gamma_2. \quad (10),$$

$$\gamma_1 \quad \gamma_2$$



$$a = c_2$$

$$p = p_2.$$

(9)-

$$[\Gamma_1^2, \Gamma_2^2] = [\gamma_1^1, \gamma_2^1] \cap [\gamma_1^2, \gamma_2^2]$$

$$G_3 = G_3(c_1, c_2) = \bigcup_{\gamma \in [\Gamma_1^2, \Gamma_2^2]} P_3(\gamma).$$



. 1

$$[\Gamma_1^2, \Gamma_2^2]$$

$G_2$

$G_3$ .

$c_2$ ,

$c_1 = 1 + i$ .

$c_2 = i$ .

(4),

$G_1, G_2, \dots, G_{s-1}$

$c_1, c_2, \dots, c_{s-1}$

$$[\Gamma_1^{s-1}, \Gamma_2^{s-1}] = \bigcap_{j=1}^{s-1} [\gamma_1^j, \gamma_2^j] \quad (12)$$

$$G_s = G_s(c_1, c_2, \dots, c_{s-1}) = \bigcup_{\gamma \in [\Gamma_1^{s-1}, \Gamma_2^{s-1}]} P_s(\gamma) \quad (s = \overline{2, N}). \quad (13)$$

$G_s$

$c_s$

$$c_s \in P_s(\gamma) \quad \Gamma_1^N \leq \gamma \leq \Gamma_2^N, \quad s = \overline{1, N}.$$

3.

(4)

$$c_s \in G_s \quad (s = \overline{1, N}), \quad G_1$$

(11),

$$G_s(c_1, c_2, \dots, c_{s-1}) \text{ ó}$$

(12)- (13)  $s = \overline{2, N}$ .

1)

(4) ó

2)

$$K(\Gamma_1^N) \cup K(\Gamma_2^N), \quad K(\gamma)$$

$$(6), \quad \Gamma_1^N, \Gamma_2^N \text{ ó}$$

(12).

(13),  $G_k \neq \emptyset$   $[\Gamma_1^N, \Gamma_2^N] \neq \emptyset$  . ,

$\gamma \in [\Gamma_1^N, \Gamma_2^N]$  :  $c_k \in P_k(\gamma)$  ( $k = \overline{1, N}$ ). , 2

(4)  $K(\gamma)$  .

$c_k \in P_k(\gamma)$ ,  $\Gamma_1^N \leq \gamma \leq \Gamma_2^N$ ,

$K(\Gamma_1^N) \cup K(\Gamma_2^N)$ ,  $K(\gamma)$  (6).

$\emptyset$  (

1- ) .

1-

1. / . . // . . . . ó 1999. ó 42, 4. ó . 7-12.
2. . . // / . . . . ó 2000. ó 31 ó . 19-23.
3. / . . // . . . . ó 1996. ó 39, 2. ó . 35-38.
4. / . . // . . . . ó 2009. ó 52, 4. ó . 73-80.
5. . ó . : . . . , 1974. ó 272 . / . . . . ó . : . . . , 1986. ó 174 .
7. . . . / . . . . ; [ . . . . . . . . . ] . ó . : . . . , 1985. ó 414 .
8. . . . / . . . . ó . . . , 2010. ó 217 .
9. . ó . : . . . , 2007. ó 294 .  $\lambda$  ó / . . . . ,
10. / . . . . ó . : . . . , 1983. ó 312 .
11. Lorentzen L. Continued fractions with applications / L. Lorentzen, H. Waadeland. ó Amsterdam: North-Holland, 1992. ó 606 p.
12. Perron O. Die Lehre von der Kettenbrühen / O. Perron. ó Stuttgart: Teubner, 1957. ó Bd. 2. ó 524 s.
13. Wall H. S. Analytic theory of continued fractions / H. S. Wall. ó New York: Van Norstrand, 1948. ó 433 p.