

**D**

1.

$f(z)$  ó  $\{z_n\}$ ,  $f(0) = 1$

$$M(r, f) = \max_{0 \leq \theta \leq 2\pi} |f(re^{i\theta})|, \quad L(r, f) = \min_{0 \leq \theta \leq 2\pi} |f(re^{i\theta})|,$$

$$n(r) = n(r, 0, f) = \sum_{|z_n| \leq r} 1, \quad N(r) = \int_0^r \frac{n(t)}{t} dt.$$

$\rho$   $f$

$$\rho = \lim_{r \rightarrow +\infty} \frac{\overline{\ln \ln M(r, f)}}{\ln r}.$$

$\varepsilon_\rho$  ó  $\rho$ ,

$$c(f) = \lim_{r \rightarrow +\infty} \frac{n(r)}{\ln M(r, f)}.$$

$$C(\rho) = \inf_{f \in \varepsilon_\rho} c(f).$$

[1] [2,3] ,

$$C(\rho) = \frac{1}{\pi} \sin \pi \rho, \quad 0 \leq \rho \leq 1; \tag{1}$$

$$\frac{|\sin \pi \theta|}{A_0(1 + \ln r)|\sin \pi \rho| + \pi} \leq C(\rho) \leq \frac{1}{\pi} |\sin \pi \rho|, \quad \rho > 1, \tag{2}$$

$A_0$  ó .

(2)

$$f_\rho(z) = \prod_{n=1}^{+\infty} \left( 1 + \frac{z}{n^\sigma} \right) \exp \left( \sum_{k=1}^q \frac{1}{k} \left( \frac{z}{-n^\sigma} \right)^k \right),$$

$$\sigma = \rho^{-1}, \quad q = [\rho], \quad \rho \notin \mathbb{N}, \quad n(r) \sim r^\rho \text{ i } \ln M(r, f) \sim \frac{\pi}{|\sin \pi \rho|} r^\rho \quad r \rightarrow +\infty.$$

[4]  $f$   $\rho > 1,$

$$C(\rho) < \frac{|\sin \pi \rho|}{\pi}.$$

$$\frac{C(\rho)}{C(\rho)} \quad \rho > 1 \quad \rho \rightarrow +\infty.$$

$$1 < \rho < +\infty$$

[5]

$$c(f) \geq A |\sin \pi \rho| \tag{3}$$

$$f \quad \rho, \quad \{z_n\} \quad \arg z = \pi.$$

[5]

(z)  $\emptyset$

$$\ln(z) = (-1)^q z^{q+1} \int_0^{+\infty} \frac{n(t, 0)}{t^{q+1}(t+z)} dt,$$

$$q = [\rho], \quad |\arg z| < \pi.$$

$$B(r, \theta) = \sup_{|\theta| < \pi} |\ln(re^{i\theta})|,$$

$$\varphi(r) = \frac{B(r, \theta)}{r^{q+1}}, \quad \varphi(r) = \frac{n(r, 0)}{r^{q+1}}.$$

$$\varphi(r) \quad r \leq 0$$

$$\varphi(r) \leq 12M\varphi(r) + \pi H^* \varphi(r) + 10 \int_0^{+\infty} \frac{\varphi(t)}{t+r} dt, \tag{4}$$

$$M\varphi(r) = \sup_{\varepsilon > 0} \frac{1}{2\varepsilon} \int_{|t-r| < \varepsilon} \varphi(t) dt$$

6

$$H^* \varphi(r) = \frac{1}{\pi} \sup_{\varepsilon > 0} \left| \int_{|t-r| > \varepsilon} \frac{\varphi(t) dt}{t-r} \right|$$

6

©

(4) [5] :

$$\left( \int_{R_n}^{+\infty} \left( \frac{B(r, f)}{r^{q+1}} \right)^p dr \right)^{1/p} \leq \frac{1}{A \sin \frac{\pi}{p}} \left( \int_{R_n}^{+\infty} \left( \frac{n(r, 0, f)}{r^{q+1}} \right)^p dr \right)^{1/p}, \tag{5}$$

$$p = \frac{1}{q+1-\rho} + \varepsilon_n, \quad R_n \rightarrow +\infty, \quad \varepsilon_n \rightarrow 0 \quad n \rightarrow +\infty \quad \acute{o} . \tag{3}$$

[6] - (4), 4 .

$$(r) \leq \pi M \varphi(r) + \pi M (H\varphi)r + \int_0^{+\infty} \frac{\varphi(t)}{t+r} dt, \tag{6}$$

$$H\varphi(r) = \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0} \int_{|t-r|>\varepsilon} \frac{\varphi(t)}{r-t} dt$$

ó .

(6) :

$$(r) \leq \pi \varphi^\perp(r) + \pi \tilde{\varphi}^\perp(r) + \int_0^{+\infty} \frac{\varphi(t)}{t+r} dt, \tag{7}$$

$$\varphi^\perp(r) = \sup_{y>0} \left| \frac{1}{\pi} \int_{-\infty}^{+\infty} \frac{y\varphi(t)dt}{(r-t)^2 + y^2} \right|$$

ó ,

$$\tilde{\varphi}^\perp(r) = \sup_{y>0} \left| \frac{1}{\pi} \int_{-\infty}^{+\infty} \frac{(r-t)\varphi(t)dt}{(r-t)^2 + y^2} \right|$$

ó .

$\|M\varphi\|_p$   $\|M(H\varphi)\|_p$   $M\varphi$   $M(H\varphi)$   $L_p$  ó  $\|\varphi_\alpha^*\|_p$   $\|\tilde{\varphi}_\alpha^*\|_p$

$$\varphi_\alpha^*(r) = \sup_{\alpha(r)} \left| \frac{1}{\pi} \int_{-\infty}^{+\infty} \frac{y\varphi(t)dt}{(x-t)^2 + y^2} \right|$$

$$\tilde{\Phi}_\alpha^*(r) = \sup_{\alpha(r)} \left| \frac{1}{\pi} \int_{-\infty}^{+\infty} \frac{(x-t)\varphi(t)dt}{(x-t)^2 + y^2} \right|$$

$$\alpha(r) = \{x + iy : |x - r| < \alpha y, \alpha > 0, y > 0\},$$

( [7, .239])

$$\|\tilde{\varphi}_\alpha^*\|_p \leq C_p \|\varphi_\alpha^*\|_p, \quad p > 1,$$

$$C_p = \operatorname{tg} \frac{\pi}{2p} \quad 1 < p \leq 2 \quad C_p = \operatorname{ctg} \frac{\pi}{2p} \quad \geq 2.$$

(3).

$$(3) \quad A = \frac{1}{4\pi},$$

$$\frac{|\sin \pi \rho|}{4\pi} \leq c(f) \leq \frac{|\sin \pi \rho|}{\pi}$$

$f \in \mathcal{E}_\rho$

[5],

$f$   $\rho > 1$ ,

$k$

2.

1.  $f \in \mathcal{E}_\rho$   $\rho > 1$   $\emptyset$ ,  $f(0) = 1$ ,

$$B(r, f) = \sup_{|\theta| < \pi} |\ln f(re)^{i\theta}|.$$

$$\overline{\lim}_{r \rightarrow +\infty} \frac{n(r)}{B(r, f)} \geq \frac{|\sin \pi \rho|}{4\pi}.$$

1.  $f \in \mathcal{E}_\rho$   $\rho > 1$   $\emptyset$ ,  $f(0) = 1$ .

$$\overline{\lim}_{r \rightarrow +\infty} \frac{N(r)}{B(r, f)} \geq \frac{\sin \pi \rho}{4\pi \rho}.$$

2.  $f \in \mathcal{E}_\rho$   $\rho > 1, f(0) = 1$   $f$

$k$ ,

$$\overline{\lim}_{r \rightarrow +\infty} \frac{n(r)}{\ln M(r, f)} \geq \frac{|\sin \pi \rho|}{4\pi k}, \quad \overline{\lim}_{r \rightarrow +\infty} \frac{\ln L(r, f)}{n(r)} \geq -\frac{4\pi k}{|\sin \pi \rho|},$$

$$\overline{\lim}_{r \rightarrow +\infty} \frac{N(r)}{\ln M(r, f)} \geq \frac{|\sin \pi \rho|}{4\pi \rho k}, \quad \overline{\lim}_{r \rightarrow +\infty} \frac{\ln L(r, f)}{N(r)} \geq -\frac{4\pi \rho k}{|\sin \pi \rho|}.$$

1. Pólya G. Bemerkungen über unendlichen Folgeundganzen Functionen/ G. Pólya // Math. Ann. ó 1923. ó Vol. 88. ó P.169-183.
2. Valiron G. Sur les fonctions entieres d'ordre nul et d'ordre fini, et en particulier les fonctions a correspondance reguliere / G.Valiron // Ann. Fac. Sci. Univ. Toulouse. ó 1913. ó Vol. 5, 3. ó P.117-257.
3. Valiron G. A propos d'un memorie de M. Polya / G.Valiron // Bull. Sci. Math. ó 1924. ó Vol. 48, 2. ó P.9-12.
4. . . // . ó 1973. ó .37, 3. ó . 603-629.
5. Shea D.F. Growth problems for a class of entire functions via singular integral estimates/ D.F.Shea, S.Wainger // Ill. J. Math. ó 1981. ó Vol. 25, 1. ó P. 41-50.
6. Abi-Khuzam F.F. On a singular integral estimate for the maximum modulus of a canonical product / F.F. Abi-Khuzam, B.Shayya // Ill. J. Math. ó 2000. ó Vol.11, 3. ó .551-555.
7. . . ( ) / . . // . . . . . ó 1987. ó . 15. ó .1976292.