

519.622

..

k -

k -

ø

k -

(ø)

ø

k -

$$\frac{dx}{dt} = f(x, t), \tag{1}$$

$f(x, t)$  -  $x$   $t$ ,

$$f_i(x, t) \in C_{x,t}^{(1,0)}(D), \tag{2}$$

$$D = \{0 < t < \infty; x \in \square^n\}.$$

(2)

$$f(x, t) \quad D$$

$x_i$   $x$

$$\|x\| < \delta$$

$$f(x, t) = A(t)x + \psi(x, t),$$

$$A(t) = \left( \frac{\partial f(x, t)}{\partial x} \right)_{|x=0}, \quad \psi(x, t)$$

$$\|x\| < \delta \tag{1}$$

$$\frac{dx}{dt} = A(t)x + \psi(x, t). \tag{3}$$

[1]

$x = 0$

$$\frac{dx}{dt} = A(t) \cdot x \tag{4}$$

ø

(3), (1)

$\psi(x, t)$

$$\|\psi(x, t)\| \leq C \|x\|^l, \quad (l > 1), \tag{5}$$

$\|x\|$

:

$$\|x\|_1 = \max_i |x_i|; \quad \|x\|_2 = \sum_{i=1}^n |x_i|; \quad \|x\|_3 = \sqrt{\sum_{i=1}^n |x_i|^2}.$$

ø

(1) k -

$$\sum_{s=0}^k \{\alpha_s x_{m+s} + h\beta_s \dot{x}_{m+s}\} = 0. \quad (6)$$

$$\sum_{s=0}^k \{\alpha_s x_{m+s} + h\beta_s f(x_{m+s}, t_{m+s})\} = 0, \quad (7)$$

$$k - \quad (6) \quad (1)$$

$$, \quad (7)$$

$$\emptyset \quad x(t) \quad (1) \quad h$$

$$f(x, t). \quad [2]$$

$$L[x(t)] = \sum_{s=0}^k \{\alpha_s x(t+s_h) + h\beta_s \dot{x}(t+s_h)\}$$

$$h \quad x(t).$$

$$\alpha_s, \beta_s \quad (7) \quad (6).$$

$$L[x(t)] = O(h^{p+1}) \quad x(t),$$

$p+1$

$$\sum_{s=0}^k \alpha_s = 0, \quad (8)$$

$$\sum_{s=0}^k \left\{ \alpha_s \frac{s^\nu}{\nu!} + \beta_s \frac{s^\nu}{(\nu-1)!} \right\} = 0, \quad \nu = 1, 2, \dots, p. \quad (9)$$

$$, \quad (6) \quad \alpha_s \quad \beta_s \quad (s = 1, 2, \dots, k)$$

(8) (9).

$$(6) \quad \alpha_k \neq 0, \quad \beta_k = 0 \quad k - \quad (4)$$

$$\sum_{s=0}^k \{\alpha_s E + h\beta_s A_{m+s}\} x_{m+s} = 0, \quad (10)$$

$$E - \quad n \times n, \quad A_s = A(t+hs). \quad \alpha_k \neq 0,$$

(10)  $\alpha_k,$

$$\sum_{s=0}^k \{a_s E + hb_s A_{m+s}\} x_{m+s} = 0,$$

$$a_k = 1, \quad b_k = 0, \quad a_s = \frac{\alpha_s}{\alpha_k}, \quad b_s = \frac{\beta_s}{\beta_k}, \quad (s = 0, 1, \dots, k-1). \quad (n \cdot k) -$$

$$z_m = (x_{m+k}, x_{m+k-1}, \dots, x_{m+1})^T. \quad (10)$$

$$z_m = B_{m-1} z_{m-1}, \quad (11)$$

$B_{m-1}$  ó  $(n \cdot k) \times (n \cdot k)$  -

$$B_{m-1} = \begin{pmatrix} D_{k-1} & D_{k-2} & \dots & D_1 & D_0 \\ E & \theta & \dots & \theta & \theta \\ \theta & E & \dots & \theta & \theta \\ \dots & \dots & \dots & \dots & \dots \\ \theta & \theta & \dots & E & \theta \end{pmatrix} \quad (12)$$

$D_i$  ó  $n$  -

$$D_i = -(a_i E + h b_i A_{m+i}), \quad (i = k-1, \dots, 0).$$

(3)

$$z_m = B_{m-1} z_{m-1} + G g_{m-1}, \quad (13)$$

$g_{m-1}$  ó  $(n \cdot k)$  -

$g_{m-1} = \text{colon}(\psi(x_{m+n-1}, t_{m+n-1}), \dots, \psi(x_m, t_m))$ ,  $G$  ó  $(n \cdot k) \times (n \cdot k)$  -

$$G = \begin{pmatrix} -hb_{k-1}E & -hb_{k-2}E & \dots & -hb_1E & -hb_0E \\ \theta & \theta & \dots & \theta & \theta \\ \dots & \dots & \dots & \dots & \dots \\ \theta & \theta & \dots & \theta & \theta \end{pmatrix}. \quad (14)$$

$h$   $I$ .

(11)

(13)  $\emptyset$  (3)

(13)

 $\emptyset$ 

(13)

$y_m(z_m, t_m)$   $\emptyset$

$$y_m(z_m, t_m) = B_{m-1} z_{m-1}. \quad (15)$$

 $V_m$ 

$h < h_{\max}$ :

$$\|y\| \leq V_m(y) \leq L\|y\|,$$

$$V_m(y_m) - V_{m-1}(y_{m-1}) \leq -(1-q)V_{m-1}(y_{m-1}), \quad (16)$$

$\|\cdot\|$  -

$$\begin{aligned} V_m(z_m) - V_{m-1}(z_{m-1}) &= V_m(y_m(z_m, t_m)) - V_{m-1}(z_{m-1}) + V_m(z_m) - V_m(y_m(z_m, t_m)) \\ &\leq -(1-q)V_{m-1}(z_{m-1}) + L\|z_m - y_m(z_m, t_m)\|. \end{aligned}$$

$$\|z_m - y_m(z_m, t_m)\|. \quad (15) \quad (13)$$

$$\|z_m - y_m(z_m, t_m)\| = \|G g_{m-1}\| \leq \|G\| \cdot \|g_{m-1}\| \leq h \sum_{j=1}^k |b_{k-j}| C_1 \|z_{m-1}\|^l \leq h \gamma_1 \|z_{m-1}\| \leq h \gamma_1 V_{m-1}(z_{m-1}),$$

$$\gamma_1 = C_1 \sum_{j=1}^k |b_{k-j}| \cdot \|z_{m-1}\|^{l-1}, \quad C_1 -$$

$$V_m(z_m) - V_{m-1}(z_{m-1}) \leq -(1-q)V_{m-1}(z_{m-1}) + hL\gamma_1 V_{m-1}(z_{m-1}).$$

$$V_m(z_m) - V_{m-1}(z_{m-1}) \leq -(1-q - hL\gamma_1)V_{m-1}(z_{m-1}).$$

$\emptyset$  (13)

$$1 - q - hL\gamma_1 > 0, \quad h < h_{\max} = \frac{1-q}{L\gamma_1}. \quad (17)$$

$$\psi(x, t) \quad (5),$$

 $\delta$  -

$\|x\| < \delta$

(17)

[3]

 $\emptyset$ 

(13)

(3)  $k$  -

(6).

(6)

 $\alpha_s \quad \beta_s \quad (s = 1, \dots, k)$ 

(8), (9)

 $\alpha_k \neq 0$  $\beta_k \neq 0$  - $k$  -

(6)

(4)

(10).

$$\alpha_k E + h\beta_k A_{m+k}, \quad m \in [0; 1; \dots] \quad (10)$$

$$x_{m+k} = -\sum_{s=0}^{k-1} (\alpha_k E + h\beta_k A_{m+k})^{-1} (\alpha_s E + h\beta_s A_{m+s}) x_{m+s}. \quad (18)$$

$$z_m = B_{m-1}^{(1)} z_{m-1}, \quad (19)$$

$z_m$  ó  $(n \cdot k)$ - ,  $B_{m-1}^{(1)}$  ó  $(n \cdot k) \times (n \cdot k)$ -

$$B_{m-1}^{(1)} = \begin{pmatrix} C_{k-1} & C_{k-2} & \dots & C_1 & C_0 \\ E & \theta & \dots & \theta & \theta \\ \theta & E & \dots & \theta & \theta \\ \dots & \dots & \dots & \dots & \dots \\ \theta & \dots & \dots & E & \theta \end{pmatrix} \quad (20)$$

$C_i$  ó  $n$ -

$$C_i = -(\alpha E + h\beta_k A_{m+k})^{-1} (\alpha_i E + h\beta_i A_{m+i}) \quad (i = k-1, \dots, 0). \quad (3)$$

$$z_m = B_{m-1}^{(1)} z_{m-1} + G_{m-1}^{(1)} g_{m-1} + G_m^{(2)} g_m, \quad (21)$$

$G_{m-1}^{(1)}$   $G_m^{(2)}$  ó  $(n \cdot k) \times (n \cdot k)$ - :

$$G_{m-1}^{(1)} = \begin{pmatrix} -h\beta_{k-1} (\alpha_k E + h\beta_k A_{m+k})^{-1} & \dots & -h\beta_0 (\alpha_k E + h\beta_k A_{m+k})^{-1} \\ \theta & \dots & \theta \\ \dots & \dots & \dots \\ \theta & \dots & \theta \end{pmatrix},$$

$$G_m^{(2)} = \begin{pmatrix} -h\beta_k (\alpha_k E + h\beta_k A_{m+k})^{-1} & \theta & \dots & \theta \\ \theta & \theta & \dots & \theta \\ \dots & \dots & \dots & \dots \\ \theta & \theta & \dots & \theta \end{pmatrix}. \quad (21) \quad [4]$$

$$\det \left( G_m^{(2)} \left( \frac{\partial g}{\partial z} \right)_{z=z_m} - E \right) \neq 0 \quad \forall z \in R^{n \cdot k}$$

$$\|G_m^{(2)} g_m(z_m, t_m) - z_m\| \rightarrow \infty, \quad \|z\| \rightarrow \infty. \quad (22)$$

$$(21) \quad \text{2.} \quad \emptyset \quad (22) \quad h \quad (21) \quad ,$$

$$\emptyset \quad (21) \quad h < h_{\max}.$$

$$y_m(z_m, t_m) \quad \emptyset$$

$$y_m(z_m, t_m) = B_{m-1}^{(1)} z_{m-1}. \quad (23)$$

$V_m$ ,

$$(16). \quad , \quad 1$$

$$V_m(z_m) - V_{m-1}(z_{m-1})$$

$$V_m(z_m) - V_{m-1}(z_{m-1}) \leq -(1-q)V_{m-1}(z_{m-1}) + L \|z_m - y_m(z_m, t_m)\|.$$

$$\|z_m - y_m(z_m, t_m)\|. \quad (23) \quad (21)$$

$$\begin{aligned} \|z_m - y_m(z_m, t_m)\| &= \|G_{m-1}^{(1)}g_{m-1} + G_m^{(2)}g_m\| \leq \|G_{m-1}^{(1)}\| \cdot \|g_{m-1}\| + \|G_m^{(2)}\| \cdot \|g_m\| \leq \\ &\leq \sum_{j=1}^k \frac{|\beta_{k-j}|}{|\beta_j| \cdot \|A_{m+k}\|} \cdot \|g_{m-1}\| + \frac{\|g_m\|}{\|A_{m+k}\|} \leq \xi_1(z_{m-1})\|z_{m-1}\| + \xi_2(z_m)\|z_m\| \leq \\ &\leq \xi_1(z_{m-1})V_{m-1}(z_{m-1}) + \xi_2(z_m)V_m(z_m), \\ \xi_1(z_{m+1}) &= \sum_{j=1}^k \frac{|\beta_{k-j}| \cdot C_1}{|\beta_j| \cdot \|A_{m+k}\|} \cdot \|z_{m-1}\|^{l-1}; \quad \xi_2(z_m) = \frac{C_2}{\|A_{m+k}\|} \cdot \|z_m\|^{l-1}, \quad C_1 \quad C_2 \end{aligned}$$

$$V_m(z_m) - V_{m-1}(z_{m-1}) \leq -(1-q)V_{m-1}(z_{m-1}) + L\xi_1 V_{m-1}(z_{m-1}) + L\xi_2 V_m(z_m).$$

$$V_m(z_m) - V_{m-1}(z_{m-1}) \leq \frac{-(1-q-L(\xi_1 + \xi_2))}{1-L\xi_2} V_{m-1}(z_{m-1}).$$

$$(21) \quad 1 - q - L(\xi_1 + \xi_2) > 0, \quad 1 - L\xi_1 > 0.$$

$$\xi_1 + \xi_2 < \frac{1-q}{L}. \tag{24}$$

$$\psi(x, t) \tag{5}$$

$$\|x\| < \delta$$

$$h < h_{\max},$$

$$(23)$$

$$(24)$$

$$[3] \quad (21)$$

1. . . . ., " ", 1967 . 472 .
2. Dahlquist G.G. Convergence and stability in the numerical integration of ordinary differential equations. "Math.Scand", 1956, vol.4 p.33-53.
3. . . . ., " ", 1971., 310 .
4. Palais R.S. Natural operations on differential forms. "Trans.Amer.Math.Soc.", 1959, vol.92, p.125-141.